


Name: Adult, Bill, G ID: ASDF0135 Session Date: 1/10/2005 Session: 1 Clinician: Clinician, Sue Clinic: South Clinic Diagnosis: Dysthymia	Alert Status: NA Most Recent Score: 78 Initial Score: 78 Change From Initial: No Reliable Change Current Distress Level: Moderate																				
Most Recent Critical Item Status: 8. Suicide - I have thoughts of ending my life. Never 11. Substance Abuse - After heavy drinking, I need a drink the next morning to get going. Frequently 26. Substance Abuse - I feel annoyed by people who criticize my drinking. Rarely 32. Substance Abuse - I have trouble at work/school because of drinking or drug use. Frequently 44. Work Violence - I feel angry enough at work/school to do something I might regret. Rarely	<table><tr><th>Subscales</th><th>Current</th><th>Outpat. Norm</th><th>Comm. Norm</th></tr><tr><td>Symptom Distress:</td><td>39</td><td>49</td><td>25</td></tr><tr><td>Interpersonal Relations:</td><td>19</td><td>20</td><td>10</td></tr><tr><td>Social Role:</td><td>20</td><td>14</td><td>10</td></tr><tr><td>Total:</td><td>78</td><td>83</td><td>45</td></tr></table>	Subscales	Current	Outpat. Norm	Comm. Norm	Symptom Distress:	39	49	25	Interpersonal Relations:	19	20	10	Social Role:	20	14	10	Total:	78	83	45
Subscales	Current	Outpat. Norm	Comm. Norm																		
Symptom Distress:	39	49	25																		
Interpersonal Relations:	19	20	10																		
Social Role:	20	14	10																		
Total:	78	83	45																		



Inpatient - Acute care settings with short stay

MH Center - Outpatient settings such as community mental health centers

MBH - Managed Behavior Health company outpatient settings

EAP - Employee Assistance Programs and students presenting at University Counseling Centers

Community Sample - Individuals randomly drawn from the community

Feedback Message:

Scores in this range are typical of persons who enter outpatient psychotherapy. It can be anticipated that a reduction of the score by 14 points or more will occur within the first eight sessions of treatment. Most patients within this score range show a reliable benefit from treatment interventions. At the end of treatment that lasts about 18 sessions, the majority of clients with a score in this range will report distress that is within the normal range of functioning.


REMINDER: THE USER IS SOLELY RESPONSIBLE FOR ANY AND ALL DECISIONS AFFECTING PATIENT CARE. THE OQ®-A IS NOT A DIAGNOSTIC TOOL AND SHOULD NOT BE USED AS SUCH. IT IS NOT A SUBSTITUTE FOR A MEDICAL OR PROFESSIONAL EVALUATION. RELIANCE ON THE OQ®-A IS AT USER'S SOLE RISK AND RESPONSIBILITY. (SEE LICENSE FOR FULL STATEMENT OF RIGHTS, RESPONSIBILITIES & DISCLAIMERS)

Name: Child, Chris ID: ABCDE051	Alert Status: NA
Session Date: 8/26/2004 Session: 1	Most Recent Score: 92
Clinician: Clinician, Jill Clinic: North Clinic	Initial Score: 92
Diagnosis: Panic Disorder	Change From Initial: No Reliable Change
Completed by: Mother	Current Distress Level: Moderately High

Most Recent Critical Item Status:	
12. Obsessive Thoughts	Frequently
20. Hallucinations	Never
21. Self Injurious Behavior	Sometimes
22. Substance Abuse	Never
28. Paranoia	Rarely
38. Delusions	Never
41. Suicidal	Sometimes
44. Hypomania	Rarely
46. Fear of Going Crazy	Rarely
51. Aggression	Rarely
58. Eating Disorder	Rarely

Subscales	Current	Output. Norm	Comm. Norm
Intrapers. Distress:	32	26.4	8.9
Somatic:	14	7.8	3.3
Interpers. Relations:	10	10.4	0.6
Social Problems:	10	6.3	0.7
Behavioral Dys.:	16	20.1	6.8
Critical Items:	10	7.7	3.0
Total:	92	78.7	23.3

Normative Comparison Groups



Graph Legend:

Residential - Primarily adolescents in residential care ranging from 90 - 180 days

Inpatient - Acute care facilities with short stay

Outpatient - Includes both public community mental health centers and private sector settings

Juvenile Justice - Adolescents presenting at intake for adjudicated offenses

Community Sample - Individuals randomly drawn from the community

Feedback Message:

Scores in this range are typical of patients who are treated in inpatient settings. The parent/patient is reporting a high degree of disturbance/dysfunction and appears to be in considerable pain. Suicidality should be monitored. Generally patients in this range show considerable improvement following outpatient and/or inpatient interventions. In outpatient settings, a reduction of the score by 13 points or more will occur within the first eight sessions of treatment.

REMINDER: THE USER IS SOLELY RESPONSIBLE FOR ANY AND ALL DECISIONS AFFECTING PATIENT CARE. THE OQ®-A IS NOT A DIAGNOSTIC TOOL AND SHOULD NOT BE USED AS SUCH. IT IS NOT A SUBSTITUTE FOR A MEDICAL OR PROFESSIONAL EVALUATION. RELIANCE ON THE OQ®-A IS AT USER'S SOLE RISK AND RESPONSIBILITY. (SEE LICENSE FOR FULL STATEMENT OF RIGHTS, RESPONSIBILITIES & DISCLAIMERS)